

username

[New Account »](#)
[Forgot Password?](#)

Aluminum Oxide


[Advanced Search »](#)

Ads by Google

[LC/MS/MS Methods](#)

BioAnalytical method development and analysis using LC/MS
www.wofelabs.com

[100 Grams Free Alumina](#)

Try 100 grams of Dynamic Adsorbents highest quality alumina for free
www.dynamicadsorbents.com

[Silicon Nanoparticles](#)

Powders and dispersions available Below 10 nm particle diameter
www.meiorum.com

[Inorganic Standards](#)

Superior Quality, Unparalleled Service
www.spexcsp.com

 Oceanography and Atmospheric Sci. ■ Atmospheric Physics

The Potential for Ozone Depletion in Solid Rocket Motor Plumes by Heterogeneous Chemistry

Authors: [M. S. Hanning-Lee](#); [P. M. Felker](#); [J. A. Syage](#); [John J. Lamb](#); [TRW SPACE AND TECHNOLOGY GROUP REDONDO BEACH CA](#)

Abstract: Rates of ozone decomposition on **aluminum oxide** (alumina) particles were measured in a flow tube reactor equipped with molecular beam sampling mass spectrometry and ultraviolet absorption spectroscopy, and in a static reaction cell equipped with ultraviolet absorption spectroscopy. Reaction probabilities η are reported for ozone on alpha-alumina, gamma-alumina, and chromatographic alumina (hydroxylated alumina), respectively, over the temperature range -60 to 200 degrees C. This work addresses the potential for stratospheric ozone depletion by launch vehicle solid rocket motor exhaust. Considering best estimates of plume particle size distributions and dispersion rates, we calculate ozone depletion profiles, for direct decomposition on alumina only. The calculated ozone holes are rather narrow. In the worst case, ozone levels are within $5 \times 10^{(-5)}$ of ambient in the center of the plume. A simple analysis of the global impact of alumina particles on 9 ozone decomposition indicates a potential steady-state daytime depletion of $<3 \times 10^{(-9)}$ at present launch rates. The laboratory measurements presented here support atmospheric modeling efforts to understand rocket plume I stratosphere interactions. The laboratory and modeling work in turn is used to define design parameters for the High Resolution Ozone ImaGing (HIROIG) spectrometer to be used to monitor ozone and chemical composition of solid rocket motor plumes.

Limitations: APPROVED FOR PUBLIC RELEASE

Description: Final rept.

Pages: 22

Report Date: AUG 1996

Report Number: A972414

 Keywords relating to this report:

- ✦ [*OZONE DEPLETION](#)
- ✦ [*ROCKET EXHAUST](#)
- ✦ [*SOLID PROPELLANT ROCKET ENGINES](#)
- ✦ [ABSORPTION SPECTRA](#)
- ✦ [ALUMINUM OXIDES](#)
- ✦ [ATMOSPHERE MODELS](#)
- ✦ [CHEMICAL COMPOSITION](#)
- ✦ [CHROMATOGRAPHS](#)
- ✦ [DECOMPOSITION](#)
- ✦ [EXHAUST PLUMES](#)
- ✦ [HETEROGENEITY](#)
- ✦ [HIGH RESOLUTION](#)
- ✦ [HYDROXYL RADICALS](#)
- ✦ [LAUNCHING](#)
- ✦ [PARTICLE SIZE](#)
- ✦ [PROBABILITY](#)
- ✦ [SPECTROMETERS](#)
- ✦ [STEADY STATE](#)
- ✦ [STRATOSPHERE](#)
- ✦ [ULTRAVIOLET SPECTROSCOPY](#)

  Adobe PDF - \$13.95

  Printed Format - \$23.95

Please check the box for the format you wish to order.

[Shipping Terms](#)
[About Electronic Delivery](#)

 [Email This Abstract](#)

[« Back to search](#)